

兔抗 CBL 多克隆抗体

中文名称：兔抗 CBL 多克隆抗体

英文名称：Anti-CBL rabbit polyclonal antibody

别名：CBL2; NSLL; C-CBL; RNF55; FRA11B

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：CBL

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

This gene is a proto-oncogene that encodes a RING finger E3 ubiquitin ligase. The encoded protein is one of the enzymes required for targeting substrates for degradation by the proteasome. This protein mediates the transfer of ubiquitin from ubiquitin conjugating enzymes (E2) to specific substrates. This protein also contains an N-terminal phosphotyrosine binding domain that allows it to interact with numerous tyrosine-phosphorylated substrates and target them for proteasome degradation. As such it functions as a negative regulator of many signal transduction pathways. This gene has been found to be mutated or translocated in many cancers including acute myeloid leukaemia. Mut

	ations in this gene are also the cause of Noonan syndrome-like disorder
Applications:	ELISA, WB
Name of antibody:	CBL
Immunogen:	Synthetic peptide of human CBL
Full name:	Cbl proto-oncogene, E3 ubiquitin protein ligase
Synonyms :	CBL2; NSLL; C-CBL; RNF55; FRA11B
SwissProt:	P22681
ELISA Recommended dilution:	2000-5000
WB Predicted band size:	100 kDa
WB Positive control:	231 cells
WB Recommended dilution:	500-2000

