

## 兔抗 CCL4 多克隆抗体

中文名称：兔抗 CCL4 多克隆抗体

英文名称： Anti-CCL4 rabbit polyclonal antibody

别名： ACT2; G-26; HC21; LAG1; LAG-1; MIP1B; SCYA2; SCYA4; MIP1B1; AT744.1;  
MIP-1-beta

抗原： CCL4

储存： 冷冻 (-20℃)

宿主： Rabbit

相关类别： 一抗

反应种属： Human, Rat

标记物： Unconjugate

克隆类型： rabbit polyclonal

### 技术规格

#### Background:

The protein encoded by this gene is a mitogen-inducible monokine and is one of the major HIV-suppressive factors produced by CD8+ T-cells. The encoded protein is secreted and has chemokinetic and inflammatory functions. Recombinant MIP-1-beta induces a dose-dependent inhibition of different strains of HIV-1, HIV-2, and simian immunodeficiency virus (SIV). The processed form MIP-1-beta(3-69) retains the abilities to induce down-modulation of surface expression of the chemokine receptor CCR5 and to inhibit the CCR5-mediated entry of HIV-1 in T-cells. MIP-1-beta(3-69) is also a ligand for CCR1 and CCR2 isoform B.

<b>Applications:</b>	ELISA, IHC
<b>Name of antibody:</b>	CCL4
<b>Immunogen:</b>	Fusion protein of human CCL4
<b>Full name:</b>	chemokine (C-C motif) ligand 4
<b>Synonyms :</b>	ACT2; G-26; HC21; LAG1; LAG-1; MIP1B; SCYA2; SCYA4; MIP1B1; AT744.1; MIP-1-beta
<b>SwissProt:</b>	P13236
<b>ELISA Recommended dilution:</b>	1000-5000
<b>IHC positive control:</b>	Human liver cancer and Human thyroid cancer
<b>IHC Recommend dilution:</b>	50-200

