

## UBE2D1 抗原（重组蛋白）

中文名称： UBE2D1 抗原（重组蛋白）

英文名称： UBE2D1 Antigen (Recombinant Protein)

别名： ubiquitin conjugating enzyme E2 D1; SFT; UBCH5; UBC4/5; UBCH5A; E2(17)KB1

相关类别： 抗原

储存： 冷冻（-20℃）

### 概述

Fusion protein corresponding to a region derived from 1-147 amino acids of human UBE2D1

### 技术规格

<b>Full name:</b>	ubiquitin conjugating enzyme E2 D1
<b>Synonyms:</b>	SFT; UBCH5; UBC4/5; UBCH5A; E2(17)KB1
<b>Swissprot:</b>	P51668
<b>Gene Accession:</b>	BC005980
<b>Purity:</b>	>85%, as determined by Coomassie blue stained SDS-PAGE
<b>Expression system:</b>	Escherichia coli
<b>Tags:</b>	His tag C-Terminus, GST tag N-Terminus
<b>Background:</b>	The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-protein ligases, or E3s. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. This enzyme is closely related to a stimulator of iron transport (SFT), and is up-regulated in hereditary hemochromatosis. It also functions in the ubiquitination of the tumor-suppressor protein p53 and

the hypoxia-inducible transcription factor HIF1alpha by interacting with the E1 ubiquitin-activating enzyme and the E3 ubiquitin-protein ligases. Two transcript variants encoding different isoforms have been found for this gene.