

## 兔抗 ZBTB44 多克隆抗体

中文名称：兔抗 ZBTB44 多克隆抗体

英文名称：Anti-ZBTB44 rabbit polyclonal antibody

别名：zinc finger and BTB domain containing 44; BTBD15; ZNF851; HSPC063

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：ZBTB44

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

**Background:**

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. Zinc finger and BTB domain-containing protein 44 (ZBTB44), also known as BTBD15, is a 570 amino acid member of the Krüppel C2H2-type zinc-finger protein family. Localized to the nucleus, ZBTB44 contains a BTB domain, also known as a POZ domain, which inhibits DNA binding and mediates homotypic and heterotypic dimerization. Characteristics of the BTB domain suggest

	t that ZBTB44 functions as a transcription regulator. Four isoforms of ZBTB44 have been identified.
<b>Applications:</b>	ELISA, IHC
<b>Name of antibody:</b>	ZBTB44
<b>Immunogen:</b>	Fusion protein of human ZBTB44
<b>Full name:</b>	zinc finger and BTB domain containing 44
<b>Synonyms:</b>	BTBD15; ZNF851; HSPC063
<b>SwissProt:</b>	Q8NCP5
<b>ELISA Recommended dilution:</b>	5000-10000
<b>IHC positive control:</b>	Human thyroid cancer and human colorectal cancer
<b>IHC Recommend dilution:</b>	30-150



