

兔抗 TRPC6 多克隆抗体

中文名称：兔抗 TRPC6 多克隆抗体

英文名称：Anti-TRPC6 rabbit polyclonal antibody

别名：TRP6; FSGS2

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：TRPC6

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

| | |
|--------------------------|---|
| Background: | The protein encoded by this gene forms a receptor-activated calcium channel in the cell membrane. The channel is activated by diacylglycerol and is thought to be under the control of a phosphatidylinositol second messenger system. Activation of this channel occurs independently of protein kinase C and is not triggered by low levels of intracellular calcium. Defects in this gene are a cause of focal segmental glomerulosclerosis 2 (FSGS2). |
| Applications: | ELISA, IHC |
| Name of antibody: | TRPC6 |
| Immunogen: | Synthetic peptide of human TRPC6 |
| Full name: | transient receptor potential cation channel, subfamily C, member 6 |
| Synonyms : | TRP6; FSGS2 |

| | |
|-----------------------------|--|
| SwissProt: | Q9Y210 |
| ELISA Recommended dilution: | 3000-10000 |
| IHC positive control: | Human liver cancer and Human lung cancer |
| IHC Recommend dilution: | 50-200 |

