

兔抗 TXNDC12 多克隆抗体

中文名称：兔抗 TXNDC12 多克隆抗体

英文名称：Anti-TXNDC12 rabbit polyclonal antibody

别名：thioredoxin domain containing 12; AG1; AGR1; ERP16; ERP18; ERP19; TLP19; hAG-1; PDIA16; hTLP19

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：TXNDC12

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	This gene encodes a member of the thioredoxin superfamily. Members of this family are characterized by a conserved active motif called the thioredoxin fold that catalyzes disulfide bond formation and isomerization. This protein localizes to the endoplasmic reticulum and has a single atypical active motif. The encoded protein is mainly involved in catalyzing native disulfide bond formation and displays activity similar to protein-disulfide isomerases. This protein may play a role in defense against endoplasmic reticulum stress. Alternate splicing results in both coding and non-coding variants.
Applications:	ELISA, IHC
Name of antibody:	TXNDC12

Immunogen:	Fusion protein of human TXNDC12
Full name:	thioredoxin domain containing 12
Synonyms:	AG1; AGR1; ERP16; ERP18; ERP19; TLP19; hAG-1; PDIA16 ; hTLP19
SwissProt:	O95881
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human colorectal cancer and Human liver cancer
IHC Recommend dilution:	50-200



