

小鼠抗 HADH 单克隆抗体

中文名称： 小鼠抗 HADH 单克隆抗体

英文名称： Anti-HADH mouse monoclonal antibody

别名： HAD; HADH1; HADHSC; HCDH; HHF4; MSCHAD; SCHAD

相关类别： 一抗

储存： 冷冻（-20℃） 避光

宿主： Mouse

抗原： HADH

反应种属： Human

标记物： Unconjugate

克隆类型： mouse monoclonal

技术规格

Background:	This gene is a member of the 3-hydroxyacyl-CoA dehydrogenase gene family. The encoded protein functions in the mitochondrial matrix to catalyze the oxidation of straight-chain 3-hydroxyacyl-CoAs as part of the beta-oxidation pathway. Its enzymatic activity is highest with medium-chain-length fatty acids. Mutations in this gene cause one form of familial hyperinsulinemic hypoglycemia. The human genome contains a related pseudogene of this gene on chromosome 15. [provided by RefSeq, May 2010].
Applications:	WB, IHC
Name of antibody:	HADH
Immunogen:	Fusion protein of human HADH
Full name:	hydroxyacyl-CoA dehydrogenase (HADH), transcript varia

	nt 2
Synonyms:	HAD; HADH1; HADHSC; HCDH; HHF4; MSCHAD; SCHAD
SwissProt:	Q16836
IHC positive control:	human liver tissue
IHC Recommend dilution:	25-100
WB Predicted band size:	34 kDa
WB Positive control:	HT29 cell lysate
WB Recommended dilution:	500-2000