

## 小鼠抗 HRAS 单克隆抗体

中文名称: 小鼠抗 HRAS 单克隆抗体

英文名称: Anti-HRAS mouse monoclonal antibody

别 名: C-BAS/HAS; C-H-RAS; C-HA-RAS1; CTLO; H-RASIDX; HAMSV; HRAS1; p21ras; RASH1

抗 原: HRAS

储 存: 冷冻 (-20℃) 避光

宿 主: Mouse

反应种属: Human, Dog, Rat, Monkey, Mouse

相关类别: 一抗

标 记 物: Unconjugate

克隆类型: mouse monoclonal

## 技术规格

Background:

This gene belongs to the Ras oncogene family, whose members are related to the transforming genes of ma mmalian sarcoma retroviruses. The products encoded by these genes function in signal transduction pathwa ys. These proteins can bind GTP and GDP, and they h ave intrinsic GTPase activity. This protein undergoes a continuous cycle of de- and re-palmitoylation, which r egulates its rapid exchange between the plasma mem brane and the Golgi apparatus. Mutations in this gene cause Costello syndrome, a disease characterized by in creased growth at the prenatal stage, growth deficienc y at the postnatal stage, predisposition to tumor form ation, mental retardation, skin and musculoskeletal abn ormalities, distinctive facial appearance and cardiovasc



## 全国订货电话 4008-723-722

|                          | ular abnormalities. Defects in this gene are implicated in a variety of cancers, including bladder cancer, follic ular thyroid cancer, and oral squamous cell carcinoma. Multiple transcript variants, which encode different isof orms, have been identified for this gene. [provided by RefSeq, Jul 2008]. |
|--------------------------|--|
| Applications:            | WB, IHC, IF  |
| Name of antibody:        | HRAS   |
| Immunogen:               | Fusion protein of human HRAS   |
| Full name:               | Harvey rat sarcoma viral oncogene homolog (HRAS), tr<br>anscript variant 1   |
| Synonyms:                | C-BAS/HAS; C-H-RAS; C-HA-RAS1; CTLO; H-RASIDX; H<br>AMSV; HRAS1; p21ras; RASH1   |
| SwissProt:               | P01112   |
| IHC positive control:    | human kidney tissue  |
| IHC Recommend dilution:  | 30-150   |
| WB Predicted band size:  | 21 kDa   |
| WB Positive control:     | HepG2, Hela, SVT2, A549, COS7, Jurkat, MDCK, PC12, MCF-7 cell lysates  |
| WB Recommended dilution: | 500-2000   |