

小鼠抗 ICAM3 单克隆抗体

中文名称: 小鼠抗 ICAM3 单克隆抗体

英文名称: Anti-ICAM3 mouse monoclonal antibody

别名: CD50; CDW50; ICAM-R

抗原: ICAM3

储存: 冷冻 (-20℃) 避光

宿主: Mouse

反应种属: Human, Dog, Monkey

相关类别: 一抗

标记物: Unconjugate

克隆类型: mouse monoclonal

技术规格

Background:	The protein encoded by this gene is a member of the intercellular adhesion molecule (ICAM) family. All ICAM proteins are type I transmembrane glycoproteins, contain 2-9 immunoglobulin-like C2-type domains, and bind to the leukocyte adhesion LFA-1 protein. This protein is constitutively and abundantly expressed by all leucocytes and may be the most important ligand for LFA-1 in the initiation of the immune response. It functions not only as an adhesion molecule, but also as a potent signalling molecule. [provided by RefSeq, Jul 2008].
Applications:	WB, IHC, IF
Name of antibody:	ICAM3
Immunogen:	Fusion protein of human ICAM3
Full name:	intercellular adhesion molecule 3 (ICAM3)

Synonyms:	CD50; CDW50; ICAM-R
SwissProt:	P32942
IHC positive control:	carcinoma of human lung tissue and adenocarcinoma of human breast tissue; human lymph node tissue and human kidney tissue
IHC Recommend dilution:	30-150
WB Predicted band size:	60 kDa
WB Positive control:	A549, COS7, Jurkat, MDCK cell lysates
WB Recommended dilution:	200-1000