

小鼠抗 NPTN 单克隆抗体

| | |
|-------|-------------------------------------|
| 中文名称: | 小鼠抗 NPTN 单克隆抗体 |
| 英文名称: | Anti-NPTN mouse monoclonal antibody |
| 别名: | GP55; GP65; np55; np65; SDFR1; SDR1 |
| 相关类别: | 一抗 |
| 储存: | 冷冻 (-20℃) 避光 |
| 宿主: | Mouse |
| 抗原: | NPTN |
| 反应种属: | Human, Dog, Rat, Monkey |
| 标记物: | Unconjugate |
| 克隆类型: | mouse monoclonal |

技术规格

| | |
|------------------------------|--|
| Background: | This gene encodes a type I transmembrane protein belonging to the Ig superfamily. The protein is believed to be involved in cell-cell interactions or cell-substrate interactions. Alternative splicing results in multiple transcript variants. [provided by RefSeq]. |
| Applications: | WB, IHC, IF |
| Name of antibody: | NPTN |
| Immunogen: | Fusion protein of human NPTN |
| Full name: | neuroplastin (NPTN), transcript variant a |
| Synonyms: | GP55; GP65; np55; np65; SDFR1; SDR1 |
| SwissProt: | Q9Y639 |
| IHC positive control: | adenocarcinoma of human ovary tissue and human pancreas tissue; carcinoma of human pancreas tissue and carcinoma of huma |

| | |
|---------------------------------|---|
| | n bladder tissue |
| IHC Recommend dilution: | 30-150 |
| WB Predicted band size: | 44 kDa |
| WB Positive control: | Human omentum tissue, Human uterus tissue, Human brain tissue, Human ovarian tissue, Human thyroid gland tissue, Human colon tissue lysates |
| WB Recommended dilution: | 500-2000 |