

## 兔抗 HSD17B6 多克隆抗体

中文名称：兔抗 HSD17B6 多克隆抗体

英文名称：Anti-HSD17B6 rabbit polyclonal antibody

别名：HSE; RODH; SDR9C6

相关类别：一抗

抗原：HSD17B6

储存：冷冻（-20℃）

宿主：Rabbit

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

|                              |   |
|------------------------------|---|
| <b>Background:</b>           | The protein encoded by this gene has both oxidoreductase and epimerase activities and is involved in androgen catabolism. The oxidoreductase activity can convert 3 alpha-diol to dihydrotestosterone, while the epimerase activity can convert androsterone to epi-androsterone. Both reactions use NAD+ as the preferred cofactor. This gene is a member of the retinol dehydrogenase family. |
| <b>Applications:</b>         | ELISA, WB, IHC  |
| <b>Name of antibody:</b>     | HSD17B6   |
| <b>Immunogen:</b>            | Fusion protein of human HSD17B6   |
| <b>Full name:</b>            | hydroxysteroid (17-beta) dehydrogenase 6  |
| <b>Synonyms :</b>            | HSE; RODH; SDR9C6   |
| <b>SwissProt:</b>            | O14756  |
| <b>ELISA Recommended dil</b> | 1000-2000   |

|                                 |                    |
|---------------------------------|--------------------|
| <b>ution:</b>                   |                    |
| <b>IHC positive control:</b>    | Human colon cancer |
| <b>IHC Recommend dilution:</b>  | 25-100             |
| <b>WB Predicted band size:</b>  | 36 kDa             |
| <b>WB Positive control:</b>     | Mouse liver tissue |
| <b>WB Recommended dilution:</b> | 200-1000           |



