

兔抗 IFT57 多克隆抗体

中文名称:	兔抗 IFT57 多克隆抗体
英文名称:	Anti-IFT57 rabbit polyclonal antibody
别名:	intraflagellar transport 57; HIPPI; MHS4R2; ESRRBL1
储存:	冷冻 (-20℃)
抗原:	IFT57
宿主:	Rabbit
反应种属:	Human, Mouse
相关类别:	一抗
标记物:	Unconjugate
克隆类型:	rabbit polyclonal

技术规格

Background:	Required for the formation of cilia. Plays an indirect role in sonic hedgehog signaling, cilia being required for all activity of the hedgehog pathway (By similarity). Has pro-apoptotic function via its interaction with HIP1, leading to recruit caspase-8 (CASP8) and trigger apoptosis. Has the ability to bind DNA sequence motif 5'-AAAGACATG-3' present in the promoter of caspase genes such as CASP1, CASP8 and CASP10, suggesting that it may act as a transcription regulator; however the relevance of such function remains unclear.
Applications:	ELISA, IHC
Name of antibody:	IFT57
Immunogen:	Fusion protein of human IFT57

Full name:	intraflagellar transport 57
Synonyms:	HIPPI; MHS4R2; ESRRBL1
SwissProt:	Q9NWB7
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human tonsil
IHC Recommend dilution:	50-300

