

兔抗 IL4 多克隆抗体

中文名称：兔抗 IL4 多克隆抗体

英文名称：Anti-IL4 rabbit polyclonal antibody

别名：BSF1; IL-4; BCGF1; BSF-1; BCGF-1

相关类别：一抗

储存：冷冻（-20℃）避光

宿主：Rabbit

抗原：IL4

反应种属：Human

标记物：Unconjugate

克隆类型：Unconjugate

技术规格

Background:	<p>The protein encoded by this gene is a pleiotropic cytokine produced by activated T cells. This cytokine is a ligand for interleukin 4 receptor. The interleukin 4 receptor also binds to IL13, which may contribute to many overlapping functions of this cytokine and IL13. STAT6, a signal transducer and activator of transcription, has been shown to play a central role in mediating the immune regulatory signal of this cytokine. This gene, IL3, IL5, IL13, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL13. This gene, IL13 and IL5 are found to be regulated coordinately by several long-range regulatory elements in an over 120 kilobase range on the chromosome. Two alternatively spliced transcript variants of this gene encoding disti</p>
--------------------	--

	nct isoforms have been reported.
Applications:	WB
Name of antibody:	IL4
Immunogen:	Synthesized peptide derived from C-terminal of human IL4.
Full name:	interleukin 4
Synonyms :	BSF1; IL-4; BCGF1; BSF-1; BCGF-1
SwissProt:	P05112
WB Predicted band size:	17 kDa
WB Positive control:	Hela cells lysate
WB Recommended dilution:	500-3000



