

## 兔抗 ITGA2 多克隆抗体

中文名称: 兔抗 ITGA2 多克隆抗体

英文名称: Anti-ITGA2 rabbit polyclonal antibody

别 名: BR; GPIa; CD49B; HPA-5; VLA-2; VLAA2

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗 原: ITGA2

反应种属: Human

标 记 物: Unconjugate

克隆类型: rabbit polyclonal


## 技术规格

| Background:       | This gene encodes the alpha subunit of a transmembrane receptor for collagens and related proteins. The encoded protein forms a heterodimer with a beta subunit and mediates the adhesion of platelets and other cell types to the extracellular matrix. Loss of the encoded protein is associated with bleeding disorder platelet-type 9. Antibodies against this protein are found in several immune disorders, including neonatal alloimmune thrombocytopenia. This gene is located adjacent to a related alpha subunit gene. Alternative splicing results in multiple transcript variants. |
|-------------------|--|
| Applications:     | ELISA, WB, IHC   |
| Name of antibody: | ITGA2  |
| Immunogen:        | Synthetic peptide of human ITGA2   |



## 全国订货电话 4008-723-722

| Full name:                | integrin, alpha 2 (CD49B, alpha 2 subunit of VLA-2 receptor) |
|---------------------------|--|
| Synonyms:                 | BR; GPIa; CD49B; HPA-5; VLA-2; VLAA2                         |
| SwissProt:                | P17301   |
| IHC positive control:     | Human gastric cancer and human breast cancer                 |
| IHC Recommend dilution:   | 25-100   |
| WB Predicted band size:   | 129 kDa  |
| WB Positive control:      | A431 cells   |
| WB Recommended dilutio n: | 200-1000   |





全国订货电话 4008-723-722

