

兔抗 LAMB1 多克隆抗体

中文名称: 兔抗 LAMB1 多克隆抗体

英文名称: Anti-LAMB1 rabbit polyclonal antibody

别名: CLM; LIS5

抗原: LAMB1

储 存: 冷冻(-20℃) 避光

宿 主: Rabbit

反应种属: Human Mouse

相关类别: 一抗

标记物: Unconjugate

克隆类型: Unconjugate

技术规格

including cell adhesion, differentiation, migration, signaling, neuri te outgrowth and metastasis. Laminins are composed of 3 non i dentical chains: laminin alpha, beta and gamma (formerly A, B1, and B2, respectively) and they form a cruciform structure consisting of 3 short arms, each formed by a different chain, and a long arm composed of all 3 chains. Each laminin chain is a multidomain protein encoded by a distinct gene. Several isoforms of

Laminins, a family of extracellular matrix glycoproteins, are the major noncollagenous constituent of basement membranes. The y have been implicated in a wide variety of biological processes

each chain have been described. Different alpha, beta and gam ma chain isomers combine to give rise to different heterotrimeri c laminin isoforms which are designated by Arabic numerals in

Background:





全国订货电话 4008-723-722

er is laminin 1. The biological functions of the different chains a nd trimer molecules are largely unknown, but some of the chains have been shown to differ with respect to their tissue distribution, presumably reflecting diverse functions in vivo. This gene encodes the beta chain isoform laminin, beta 1. The beta 1 chain has 7 structurally distinct domains which it shares with other beta chain isomers. The C-terminal helical region containing domains I and II are separated by domain alpha, domains III and V contain several EGF-like repeats, and domains IV and VI have a globular conformation. Laminin, beta 1 is expressed in most tissues that produce basement membranes, and is one of the 3 chains constituting laminin 1, the first laminin isolated from Enge libreth-Holm-Swarm (EHS) tumor. A sequence in the beta 1 chain that is involved in cell attachment, chemotaxis, and binding to the laminin receptor was identified and shown to have the capacity to inhibit metastasis. Applications: WB, IHC, IF Name of antibody: LAMB1 Immunogen: Synthesized peptide derived from C-terminal of human LAMB1. Full name: laminin, beta 1 Synonyms: CLM; LIS5 SwissProt: P07942 IHC positive control: Human liver carcinoma tissue THC Recommend dilution: 198 kDa WB Predicted band size 198 kDa WB Positive control: HepG2 cells lysate WB Recommended dilution: 100-500		the order of their discovery, i.e. alpha1beta1gamma1 heterotrim
ns have been shown to differ with respect to their tissue distrib ution, presumably reflecting diverse functions in vivo. This gene encodes the beta chain isoform laminin, beta 1. The beta 1 chain has 7 structurally distinct domains which it shares with other beta chain isomers. The C-terminal helical region containing do mains I and II are separated by domain alpha, domains III and V contain several EGF-like repeats, and domains IV and VI have a globular conformation. Laminin, beta 1 is expressed in most tissues that produce basement membranes, and is one of the 3 chains constituting laminin 1, the first laminin isolated from Enge libreth-Holm-Swarm (EHS) tumor. A sequence in the beta 1 chain that is involved in cell attachment, chemotaxis, and binding to the laminin receptor was identified and shown to have the capacity to inhibit metastasis. Applications: WB, IHC, IF Name of antibody: LAMB1 Immunogen: Synthesized peptide derived from C-terminal of human LAMB1. Full name: laminin, beta 1 Synonyms: CLM; LIS5 SwissProt: P07942 IHC positive control: Human liver carcinoma tissue WB Predicted band size: 198 kDa WB Predicted band size: 198 kDa WB Positive control: HepG2 cells lysate WB Recommended dilution: HepG2 cells lysate WB Recommended dilution: Hela cells		er is laminin 1. The biological functions of the different chains a
ution, presumably reflecting diverse functions in vivo. This gene encodes the beta chain isoform laminin, beta 1. The beta 1 chain has 7 structurally distinct domains which it shares with other beta chain isomers. The C-terminal helical region containing domains I and II are separated by domain alpha, domains III and V contain several EGF-like repeats, and domains IV and VI have a globular conformation. Laminin, beta 1 is expressed in most ti ssues that produce basement membranes, and is one of the 3 chains constituting laminin 1, the first laminin isolated from Enge lbreth-Holm-Swarm (EHS) tumor. A sequence in the beta 1 chain that is involved in cell attachment, chemotaxis, and binding to the laminin receptor was identified and shown to have the capacity to inhibit metastasis. Applications: WB, IHC, IF Name of antibody: LAMB1 Immunogen: Synthesized peptide derived from C-terminal of human LAMB1. Full name: laminin, beta 1 Synonyms: CLM; LIS5 SwissProt: P07942 HC positive control: Human liver carcinoma tissue 198 kDa WB Predicted band size 198 kDa WB Positive control: HepG2 cells lysate WB Recommended dilution: IF positive control: Hela cells		nd trimer molecules are largely unknown, but some of the chai
encodes the beta chain isoform laminin, beta 1. The beta 1 chain has 7 structurally distinct domains which it shares with other beta chain isomers. The C-terminal helical region containing domains I and II are separated by domain alpha, domains III and V contain several EGF-like repeats, and domains IV and VI have a globular conformation. Laminin, beta 1 is expressed in most tissues that produce basement membranes, and is one of the 3 chains constituting laminin 1, the first laminin isolated from Engel Ibreth-Holm-Swarm (EHS) tumor. A sequence in the beta 1 chain that is involved in cell attachment, chemotaxis, and binding to the laminin receptor was identified and shown to have the capacity to inhibit metastasis. Applications: WB, IHC, IF Name of antibody: LAMB1 Immunogen: Synthesized peptide derived from C-terminal of human LAMB1. Full name: laminin, beta 1 Synonyms: CLM; LIS5 SwissProt: P07942 Human liver carcinoma tissue IHC positive control: Human liver carcinoma tissue 198 kDa : WB Predicted band size : WB Positive control: HepG2 cells lysate WB Recommended dilution: 500-3000 IF positive control: Hela cells		ns have been shown to differ with respect to their tissue distrib
n has 7 structurally distinct domains which it shares with other beta chain isomers. The C-terminal helical region containing do mains I and II are separated by domain alpha, domains III and V contain several EGF-like repeats, and domains IV and VI have a globular conformation. Laminin, beta 1 is expressed in most ti ssues that produce basement membranes, and is one of the 3 c hains constituting laminin 1, the first laminin isolated from Enge lbreth-Holm-Swarm (EHS) tumor. A sequence in the beta 1 chain that is involved in cell attachment, chemotaxis, and binding to the laminin receptor was identified and shown to have the capacity to inhibit metastasis. Applications: WB, IHC, IF Name of antibody: LAMB1 Immunogen: Synthesized peptide derived from C-terminal of human LAMB1. Full name: laminin, beta 1 Synonyms: CLM; LIS5 SwissProt: P07942 HUC positive control: Human liver carcinoma tissue 198 kDa 198 kDa WB Predicted band size WB Positive control: HepG2 cells lysate WB Recommended dilution: WB Recommended dilution: Full positive control: HepG2 cells lysate WB Recommended dilution: Hela cells		ution, presumably reflecting diverse functions in vivo. This gene
beta chain isomers. The C-terminal helical region containing do mains I and II are separated by domain alpha, domains II and V contain several EGF-like repeats, and domains IV and VI have a globular conformation. Laminin, beta 1 is expressed in most ti ssues that produce basement membranes, and is one of the 3 c hains constituting laminin 1, the first laminin isolated from Enge libreth-Holm-Swarm (EHS) tumor. A sequence in the beta 1 chain that is involved in cell attachment, chemotaxis, and binding to the laminin receptor was identified and shown to have the capacity to inhibit metastasis. Applications: WB, IHC, IF Name of antibody: LAMB1 Immunogen: Synthesized peptide derived from C-terminal of human LAMB1. Full name: laminin, beta 1 Synonyms: CLM; LIS5 SwissProt: P07942 IHC positive control: Human liver carcinoma tissue IHC Recommend dilution: n: WB Predicted band size 198 kDa 198 kDa 198 kDa 198 Recommended dilution: WB Recommended dilution: HepG2 cells lysate WB Recommended dilution: Hela cells		encodes the beta chain isoform laminin, beta 1. The beta 1 chai
mains I and II are separated by domain alpha, domains III and V contain several EGF-like repeats, and domains IV and VI have a globular conformation. Laminin, beta 1 is expressed in most ti ssues that produce basement membranes, and is one of the 3 c hains constituting laminin 1, the first laminin isolated from Enge lbreth-Holm-Swarm (EHS) tumor. A sequence in the beta 1 chai n that is involved in cell attachment, chemotaxis, and binding t o the laminin receptor was identified and shown to have the ca pacity to inhibit metastasis. Applications: WB, IHC, IF Name of antibody: LAMB1 Immunogen: Synthesized peptide derived from C-terminal of human LAMB1. Full name: laminin, beta 1 Synonyms: CLM; LIS5 SwissProt: P07942 IHC positive control: Human liver carcinoma tissue IHC Recommend dilution: 50-100 WB Predicted band size : 198 kDa		n has 7 structurally distinct domains which it shares with other
V contain several EGF-like repeats, and domains IV and VI have a globular conformation. Laminin, beta 1 is expressed in most ti ssues that produce basement membranes, and is one of the 3 c hains constituting laminin 1, the first laminin isolated from Enge lbreth-Holm-Swarm (EHS) tumor. A sequence in the beta 1 chai n that is involved in cell attachment, chemotaxis, and binding t o the laminin receptor was identified and shown to have the ca pacity to inhibit metastasis. Applications: WB, IHC, IF Name of antibody: LAMB1 Immunogen: Synthesized peptide derived from C-terminal of human LAMB1. Full name: laminin, beta 1 Synonyms: CLM; LIS5 SwissProt: P07942 IHC positive control: Human liver carcinoma tissue IHC Recommend dilutio n: WB Predicted band size : WB Positive control: HepG2 cells lysate WB Recommended dilution: 500-3000 IF positive control: Hela cells		beta chain isomers. The C-terminal helical region containing do
a globular conformation. Laminin, beta 1 is expressed in most ti ssues that produce basement membranes, and is one of the 3 c hains constituting laminin 1, the first laminin isolated from Enge lbreth-Holm-Swarm (EHS) tumor. A sequence in the beta 1 chai n that is involved in cell attachment, chemotaxis, and binding t o the laminin receptor was identified and shown to have the ca pacity to inhibit metastasis. Applications: WB, IHC, IF Name of antibody: LAMB1 Immunogen: Synthesized peptide derived from C-terminal of human LAMB1. Full name: laminin, beta 1 Synonyms: CLM; LIS5 SwissProt: P07942 IHC positive control: Human liver carcinoma tissue IHC Recommend dilutio n: WB Predicted band size : WB Positive control: HepG2 cells lysate WB Recommended dilution: 500-3000 IF positive control: Hela cells		mains I and II are separated by domain alpha, domains III and
ssues that produce basement membranes, and is one of the 3 c hains constituting laminin 1, the first laminin isolated from Enge lbreth-Holm-Swarm (EHS) tumor. A sequence in the beta 1 chai n that is involved in cell attachment, chemotaxis, and binding t o the laminin receptor was identified and shown to have the ca pacity to inhibit metastasis. Applications: WB, IHC, IF Name of antibody: LAMB1 Immunogen: Synthesized peptide derived from C-terminal of human LAMB1. Full name: laminin, beta 1 Synonyms: CLM; LIS5 SwissProt: P07942 IHC positive control: Human liver carcinoma tissue IHC Recommend dilutio n: WB Predicted band size : WB Positive control: HepG2 cells lysate WB Recommended dilut ion: IF positive control: Hela cells		V contain several EGF-like repeats, and domains IV and VI have
hains constituting laminin 1, the first laminin isolated from Enge lbreth-Holm-Swarm (EHS) tumor. A sequence in the beta 1 chai n that is involved in cell attachment, chemotaxis, and binding t o the laminin receptor was identified and shown to have the capacity to inhibit metastasis. Applications: WB, IHC, IF Name of antibody: LAMB1 Immunogen: Synthesized peptide derived from C-terminal of human LAMB1. Full name: laminin, beta 1 Synonyms: CLM; LIS5 SwissProt: P07942 IHC positive control: Human liver carcinoma tissue IHC Recommend dilution: WB Predicted band size: WB Positive control: HepG2 cells lysate WB Recommended dilution: WB Pecommended dilution: Hela cells		
Ibreth-Holm-Swarm (EHS) tumor. A sequence in the beta 1 chain that is involved in cell attachment, chemotaxis, and binding to the laminin receptor was identified and shown to have the capacity to inhibit metastasis. Applications: WB, IHC, IF Name of antibody: LAMB1 Immunogen: Synthesized peptide derived from C-terminal of human LAMB1. Full name: laminin, beta 1 Synonyms: CLM; LIS5 SwissProt: P07942 IHC positive control: Human liver carcinoma tissue IHC Recommend dilution: 50-100 WB Predicted band size: 198 kDa : WB Positive control: HepG2 cells lysate WB Recommended dilution: 500-3000 IF positive control: Hela cells		·
n that is involved in cell attachment, chemotaxis, and binding to the laminin receptor was identified and shown to have the capacity to inhibit metastasis. Applications: WB, IHC, IF Name of antibody: LAMB1 Immunogen: Synthesized peptide derived from C-terminal of human LAMB1. Full name: laminin, beta 1 Synonyms: CLM; LIS5 SwissProt: P07942 IHC positive control: Human liver carcinoma tissue IHC Recommend dilution: 50-100 n: WB Predicted band size: 198 kDa : WB Positive control: HepG2 cells lysate WB Recommended dilution: 500-3000 IF positive control: Hela cells		
o the laminin receptor was identified and shown to have the ca pacity to inhibit metastasis. Applications: WB, IHC, IF Name of antibody: LAMB1 Immunogen: Synthesized peptide derived from C-terminal of human LAMB1. Full name: laminin, beta 1 Synonyms: CLM; LIS5 SwissProt: P07942 IHC positive control: Human liver carcinoma tissue IHC Recommend dilution: n: WB Predicted band size: : WB Positive control: HepG2 cells lysate WB Recommended dilution: ion: IF positive control: Hela cells		·
Applications: WB, IHC, IF Name of antibody: LAMB1 Immunogen: Synthesized peptide derived from C-terminal of human LAMB1. Full name: laminin, beta 1 Synonyms: CLM; LIS5 SwissProt: P07942 IHC positive control: Human liver carcinoma tissue IHC Recommend dilutio n: WB Predicted band size : WB Positive control: HepG2 cells lysate WB Recommended dilution: 500-3000 IF positive control: Hela cells		_
Applications:WB, IHC, IFName of antibody:LAMB1Immunogen:Synthesized peptide derived from C-terminal of human LAMB1.Full name:laminin, beta 1Synonyms:CLM; LIS5SwissProt:P07942IHC positive control:Human liver carcinoma tissueIHC Recommend dilution:50-100n:198 kDaWB Predicted band size:198 kDaWB Recommended dilution:500-3000IF positive control:Hela cells		
Name of antibody:LAMB1Immunogen:Synthesized peptide derived from C-terminal of human LAMB1.Full name:laminin, beta 1Synonyms:CLM; LIS5SwissProt:P07942IHC positive control:Human liver carcinoma tissueIHC Recommend dilutio n:50-100WB Predicted band size :198 kDaWB Positive control:HepG2 cells lysateWB Recommended dilution:500-3000IF positive control:Hela cells		
Immunogen:Synthesized peptide derived from C-terminal of human LAMB1.Full name:laminin, beta 1Synonyms:CLM; LIS5SwissProt:P07942IHC positive control:Human liver carcinoma tissueIHC Recommend dilutio n:50-100WB Predicted band size :198 kDaWB Positive control:HepG2 cells lysateWB Recommended dilution:500-3000IF positive control:Hela cells		
Full name: laminin, beta 1 Synonyms: CLM; LIS5 SwissProt: P07942 IHC positive control: Human liver carcinoma tissue IHC Recommend dilutio n: WB Predicted band size : WB Positive control: HepG2 cells lysate WB Recommended dilutio ton: IF positive control: Hela cells	-	
Synonyms: CLM; LIS5 SwissProt: P07942 IHC positive control: Human liver carcinoma tissue IHC Recommend dilution: 50-100 WB Predicted band size: 198 kDa WB Positive control: HepG2 cells lysate WB Recommended dilution: 500-3000 IF positive control: Hela cells		
SwissProt:P07942IHC positive control:Human liver carcinoma tissueIHC Recommend dilutio n:50-100WB Predicted band size :198 kDa:WB Positive control:HepG2 cells lysateWB Recommended dilution:500-3000IF positive control:Hela cells	Full name:	laminin, beta 1
IHC positive control: Human liver carcinoma tissue IHC Recommend dilution: 50-100 WB Predicted band size: 198 kDa WB Positive control: HepG2 cells lysate WB Recommended dilution: 500-3000 IF positive control: Hela cells	Synonyms:	CLM; LIS5
IHC Recommend dilutio n: WB Predicted band size : WB Positive control: WB Recommended dilut ion: IF positive control: Hela cells	SwissProt:	
n: WB Predicted band size : 198 kDa : WB Positive control: HepG2 cells lysate WB Recommended dilut ion: IF positive control: Hela cells	IHC positive control:	Human liver carcinoma tissue
: 198 kDa WB Positive control: HepG2 cells lysate WB Recommended dilut ion: 500-3000 IF positive control: Hela cells		50-100
WB Recommended dilut ion: IF positive control: Hela cells	WB Predicted band size :	198 kDa
ion: IF positive control: Hela cells	WB Positive control:	HepG2 cells lysate
		500-3000
IF Recommend dilution: 100-500	IF positive control:	Hela cells
	IF Recommend dilution:	100-500



全国订货电话 4008-723-722







全国订货电话 4008-723-722

