

兔抗 CTAG1A/B 多克隆抗体

中文名称: 兔抗 CTAG1A/B 多克隆抗体

英文名称: Anti-CTAG1A/B rabbit polyclonal antibody

别名: CTAG, CTAG1, ESO1, LAGE2, LAGE2A/CTAG1B

相关类别: 一抗

储存: 冷冻 (-20℃)

宿主: Rabbit

抗原: CTAG1A/B

反应种属: Human

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:	The protein encoded by this gene is an antigen that is overexpressed in many cancers but that is also expressed in normal testis. This gene is found in a duplicated region of the X-chromosome and therefore has a neighboring gene of identical sequence. Expressed in testis and ovary and in a wide variety of cancers. Detected in uterine myometrium . Expressed from 18 weeks until birth in human fetal testis . In the adult testis, is strongly expressed in spermatogonia and in primary spermatocytes, but not in post-meiotic cells or in testicular somatic cells (at protein level).
Applications:	ELISA, IHC
Name of antibody:	CTAG1A/B
Immunogen:	Synthetic peptide of human CTAG1A/B

Full name:	Cancer/testis antigen 1
Synonyms:	CTAG, CTAG1, ESO1, LAGE2, LAGE2A/CTAG1B
SwissProt:	P78358/P78358
IHC positive control:	Human gastric cancer and human thyroid cancer
IHC Recommend dilution:	25-100

