

兔抗 CXCL14 多克隆抗体

中文名称：兔抗 CXCL14 多克隆抗体

英文名称： Anti-CXCL14 rabbit polyclonal antibody

别名： KEC; KS1; BMAC; BRAK; NJAC; MIP2G; MIP-2g; SCYB14

抗原： CXCL14

储存： 冷冻（-20℃）

宿主： Rabbit

相关类别： 一抗

反应种属： Human

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:	This gene belongs to the cytokine gene family which encode secreted proteins involved in immunoregulatory and inflammatory processes. The protein encoded by this gene is structurally related to the CXC (Cys-X-Cys) subfamily of cytokines. Members of this subfamily are characterized by two cysteines separated by a single amino acid. This cytokine displays chemotactic activity for monocytes but not for lymphocytes, dendritic cells, neutrophils or macrophages. It has been implicated that this cytokine is involved in the homeostasis of monocyte-derived macrophages rather than in inflammation.
Applications:	ELISA, IHC

Name of antibody:	CXCL14
Immunogen:	Fusion protein of human CXCL14
Full name:	chemokine (C-X-C motif) ligand 14
Synonyms :	KEC; KS1; BMAC; BRAK; NJAC; MIP2G; MIP-2g; SCYB14
SwissProt:	O95715
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human thyroid cancer
IHC Recommend dilution:	50-200

