

CST4 抗原（重组蛋白）

中文名称：CST4 抗原（重组蛋白）

英文名称：CST4 Antigen (Recombinant Protein)

储 存： 冷冻（-20℃）

相关类别： 抗原

概 述：

Fusion protein corresponding to a region derived from 21-141 amino acids of human CST4

技术规格：

Full name:	cystatin S
Swissprot:	P01036
Gene Accession:	BC065714
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE
Expression system:	Escherichia coli
Tags:	His tag C-Terminus, GST tag N-Terminus
Background:	The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins and the kininogens. The type 2 cystatin proteins are a class of cysteine proteinase inhibitors found in a variety of human fluids and secretions. The cystatin locus on chromosome 20 contains the majority of the type 2 cystatin genes and pseudogenes. This gene is located in the cystatin locus and encodes a type 2 salivary cysteine peptidase inhibitor. The protein is an S-type cystatin, based on its high level of expression in saliva, tears and seminal plasma. The specific

role in these fluids is unclear but antibacterial and antiviral activity is present, consistent with a protective function.