

## 兔抗 MAP3K8(Ab-400) 多克隆抗体

- 中文名称: 兔抗 MAP3K8(Ab-400) 多克隆抗体
- 英文名称: Anti-MAP3K8(Ab-400) rabbit polyclonal antibody
- 别名: COT; EST; ESTF; TPL2; MEKK8; Tpl-2; c-COT
- 相关类别: 一抗
- 储存: 冷冻 (-20℃) 避光
- 宿主: Rabbit
- 抗原: MAP3K8(Ab-400)
- 反应种属: Human
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

### 技术规格

<b>Background:</b>	This gene is an oncogene that encodes a member of the serine/threonine protein kinase family. The encoded protein localizes to the cytoplasm and can activate both the MAP kinase and JNK kinase pathways. This protein was shown to activate I $\kappa$ B kinases, and thus induce the nuclear production of NF- $\kappa$ B. This protein was also found to promote the production of TNF- $\alpha$ and IL-2 during T lymphocyte activation. This gene may also utilize a downstream in-frame translation start codon, and thus produce an isoform containing a shorter N-terminus.
<b>Applications:</b>	WB, IHC

<b>Name of antibody:</b>	MAP3K8(Ab-400)
<b>Immunogen:</b>	Synthesized non-phosphopeptide derived from human MAP3K8 around the phosphorylation site of serine 400 (C-Q-S(p)-L-D).
<b>Full name:</b>	mitogen-activated protein kinase kinase kinase 8
<b>Synonyms :</b>	COT; EST; ESTF; TPL2; MEKK8; Tpl-2; c-COT
<b>SwissProt:</b>	P41279
<b>IHC positive control:</b>	Human brain tissue
<b>IHC Recommend dilution:</b>	50-100
<b>WB Predicted band size:</b>	53 kDa
<b>WB Positive control:</b>	COLO cells treated with Insulin (0.01U/ml, 15mins) lysate
<b>WB Recommended dilution:</b>	500-3000



