

PSMC2 抗原（重组蛋白）

中文名称： PSMC2 抗原（重组蛋白）

英文名称： PSMC2 Antigen (Recombinant Protein)

别名： proteasome (prosome, macropain) 26S subunit, ATPase, 2; S7; MSS1; Nbla10058

储存： 冷冻（-20℃）

相关类别： 抗原

概述

Fusion protein corresponding to N terminal 250 amino acids of human PSMC2

技术规格

Full name:	proteasome (prosome, macropain) 26S subunit, ATPase, 2
Synonyms:	S7; MSS1; Nbla10058
Swissprot:	P35998
Gene Accession:	BC002589
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE
Expression system:	Escherichia coli
Tags:	His tag C-Terminus, GST tag N-Terminus
Background:	The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is

the processing of class I MHC peptides. This gene encodes one of the ATPase subunits, a member of the triple-A family of ATPases which have a chaperone-like activity.