

兔抗 ELAC2 多克隆抗体

中文名称：兔抗 ELAC2 多克隆抗体

英文名称： Anti-ELAC2 rabbit polyclonal antibody

别名： ELC2; HPC2; COXPD17

相关类别： 一抗

储存： 冷冻（-20℃）

抗原： ELAC2

宿主： Rabbit

反应种属： Human, Mouse, Rat

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

| | |
|--------------------------|--|
| Background: | The protein encoded by this gene has a C-terminal domain with tRNA 3' processing endoribonuclease activity, which catalyzes the removal of the 3' trailer from precursor tRNAs. The protein also interacts with activated Smad family member 2 (Smad2) and its nuclear partner forkhead box H1 (also known as FAST-1), and reduced expression can suppress transforming growth factor-beta induced growth arrest. Mutations in this gene result in an increased risk of prostate cancer. Multiple transcript variants encoding different isoforms have been found for this gene. |
| Applications: | ELISA, WB, IHC |
| Name of antibody: | ELAC2 |
| Immunogen: | Fusion protein of human ELAC2 |

| | |
|---------------------------------|---|
| Full name: | elaC ribonuclease Z 2 |
| Synonyms: | ELC2; HPC2; COXPD17 |
| SwissProt: | Q9BQ52 |
| IHC positive control: | Human gastric cancer and human liver cancer |
| IHC Recommend dilution: | 50-200 |
| WB Predicted band size: | 92 kDa |
| WB Positive control: | LoVo and Hela cells |
| WB Recommended dilution: | 200-1000 |



