

## 兔抗 H2AC12 多克隆抗体

中文名称：兔抗 H2AC12 多克隆抗体

英文名称： Anti-H2AC12 rabbit polyclonal antibody

别名： H2A clustered histone 12; H2AH; H2A/S; H2AFALii; H

相关类别： 一抗

抗原： H2AC12

储存： 冷冻 (-20℃)

宿主： Rabbit

反应种属： Human, Mouse

标记物： Unconjugate

克隆类型： rabbit polyclonal

### 技术规格

#### Background:

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H2A family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is f

	ound in the histone microcluster on chromosome 6p21.33. [provided by RefSeq, Aug 2015]
<b>Applications:</b>	ELISA, WB, IHC
<b>Name of antibody:</b>	H2AC12
<b>Immunogen:</b>	Fusion protein of human H2AC12
<b>Full name:</b>	H2A clustered histone 12
<b>Synonyms:</b>	H2AH; H2A/S; H2AFALii; HIST1H2AH; dJ86C11.1
<b>SwissProt:</b>	Q96KK5
<b>IHC positive control:</b>	Human colon cancer and Human liver cancer
<b>IHC Recommend dilution:</b>	50-200
<b>WB Predicted band size:</b>	14 kDa
<b>WB Positive control:</b>	Mouse liver tissue, HepG2, 293T, Hela, Raji, A375 and K562 cells
<b>WB Recommended dilution:</b>	500-2000



