

兔抗 EPCAM 多克隆抗体

中文名称：兔抗 EPCAM 多克隆抗体

英文名称：Anti-EPCAM rabbit polyclonal antibody

别名：epithelial cell adhesion molecule; ESA; KSA; M4S1; MK-1; DIAR5; EGP-2; EGP40; KS1/4; MIC18; TROP1; EGP314; HNPCC8; TACSTD1

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：EPCAM

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	This gene encodes a carcinoma-associated antigen and is a member of a family that includes at least two type I membrane proteins. This antigen is expressed on most normal epithelial cells and gastrointestinal carcinomas and functions as a homotypic calcium-independent cell adhesion molecule. The antigen is being used as a target for immunotherapy treatment of human carcinomas. Mutations in this gene result in congenital tufting enteropathy.
Applications:	ELISA, WB, IHC
Name of antibody:	EPCAM
Immunogen:	Fusion protein of human EPCAM

Full name:	epithelial cell adhesion molecule
Synonyms:	ESA; KSA; M4S1; MK-1; DIAR5; EGP-2; EGP40; KS1/4; MIC18; TROP1; EGP314; HNPCC8; TACSTD1
SwissProt:	P16422
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human liver cancer and Human prostate cancer
IHC Recommend dilution:	50-200
WB Predicted band size:	35 kDa
WB Positive control:	HT-29 cell lysate
WB Recommended dilution:	1000-5000



