

兔抗 MUC15 多克隆抗体

| | |
|-------|---------------------------------------|
| 中文名称: | 兔抗 MUC15 多克隆抗体 |
| 英文名称: | Anti-MUC15 rabbit polyclonal antibody |
| 别名: | PAS3; MUC-15; PASIII |
| 抗原: | MUC15 |
| 储存: | 冷冻 (-20℃) |
| 宿主: | Rabbit |
| 相关类别: | 一抗 |
| 反应种属: | Human |
| 标记物: | Unconjugate |
| 克隆类型: | rabbit polyclonal |

技术规格

| | |
|--------------------------------|--|
| Background: | Mucins are a family of high molecular weight, heavily glycosylated proteins (glycoconjugates) produced by epithelial tissues in most metazoans. Mucin 15 is involved in cell adhesion to the extracellular matrix. It is primarily expressed in prostate, testis, lung, spleen, thymus, ovary, small intestine, colon, bone marrow, lymph node, lung and peripheral blood leukocytes. Mucin 15 is a highly glycosylated protein. |
| Applications: | ELISA, IHC |
| Name of antibody: | MUC15 |
| Immunogen: | Fusion protein of human MUC15 |
| Full name: | mucin 15, cell surface associated |
| Synonyms : | PAS3; MUC-15; PASIII |
| SwissProt: | Q8N387 |
| ELISA Recommended dilut | 2000-5000 |

| | |
|-------------------------|--|
| ion: | |
| IHC positive control: | Human thyroid cancer and Human lung cancer |
| IHC Recommend dilution: | 25-100 |

