

兔抗 FCAR 多克隆抗体

中文名称: 兔抗 FCAR 多克隆抗体

英文名称: Anti-FCAR rabbit polyclonal antibody

别 名: CD89

相关类别: 一抗

抗原: FCAR

储 存: 冷冻(-20℃)

宿 主: Rabbit

反应种属: Human

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:


ne superfamily and encodes a receptor for the Fc region of IgA. The receptor is a transmembrane gl ycoprotein present on the surface of myeloid linea ge cells such as neutrophils, monocytes, macropha ges, and eosinophils, where it mediates immunolo gic responses to pathogens. It interacts with IgA-o psonized targets and triggers several immunologic defense processes, including phagocytosis, antibod y-dependent cell-mediated cytotoxicity, and stimul ation of the release of inflammatory mediators. M ultiple alternatively spliced transcript variants enco ding different isoforms have been described for th

This gene is a member of the immunoglobulin ge



全国订货电话 4008-723-722

	is gene.
Applications:	ELISA, WB, IHC
Name of antibody:	FCAR
Immunogen:	Fusion protein of human FCAR
Full name:	Immunoglobulin alpha Fc receptor
Synonyms:	CD89
SwissProt:	P24071
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human liver cancer and Human prostate cancer
IHC Recommend dilution:	25-100
WB Predicted band size:	32 kDa
WB Positive control:	293T cells
WB Recommended dilution:	500-2000





全国订货电话 4008-723-722

