

## 小鼠抗 KRT7 单克隆抗体

中文名称: 小鼠抗 KRT7 单克隆抗体

英文名称: Anti-KRT7 mouse monoclonal antibody

别名: CK7; K2C7; K7; SCL

相关类别: 一抗

储存: 冷冻 (-20℃) 避光

宿主: Mouse

抗原: KRT7

反应种属: Human

标记物: Unconjugate

克隆类型: mouse monoclonal

### 技术规格

|                          |  |
|--------------------------|--|
| <b>Background:</b>       | The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins which are arranged in pairs of heterotypic keratin chains coexpressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is specifically expressed in the simple epithelia lining the cavities of the internal organs and in the gland ducts and blood vessels. The genes encoding the type II cytokeratins are clustered in a region of chromosome 12q12-q13. Alternative splicing may result in several transcript variants; however, not all variants have been fully described. [provided by RefSeq, Jul 2008]. |
| <b>Applications:</b>     | WB, IHC, FC  |
| <b>Name of antibody:</b> | KRT7   |

|                                 |  |
|---------------------------------|--|
| <b>Immunogen:</b>               | Fusion protein of human KRT7                           |
| <b>Full name:</b>               | keratin 7, type II (KRT7)                              |
| <b>Synonyms:</b>                | CK7; K2C7; K7; SCL                                     |
| <b>SwissProt:</b>               | P08729   |
| <b>IHC positive control:</b>    | human kidney tissue and carcinoma of human lung tissue |
| <b>IHC Recommend dilution:</b>  | 30-150   |
| <b>WB Predicted band size:</b>  | 51 kDa   |
| <b>WB Positive control:</b>     | A549 cell lysate                                       |
| <b>WB Recommended dilution:</b> | 500-2000   |







