

兔抗 RPL17 多克隆抗体

中文名称: 兔抗 RPL17 多克隆抗体

英文名称: Anti-RPL17 rabbit polyclonal antibody

别 名: ribosomal protein L17; L17; PD-1; RPL23

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗 原: RPL17

反应种属: Human, Mouse, Rat

标 记 物: Unconjugate

克隆类型: rabbit polyclonal

技术规格


Background:

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L22P fam ily of ribosomal proteins. It is located in the cytoplasm. This gene has been referred to as rpL23 because the en coded protein shares amino acid identity with ribosomal protein L23 from Halobacterium marismortui; however, it s official symbol is RPL17. As is typical for genes encoding ribosomal proteins, there are multiple processed pseu dogenes of this gene dispersed through the genome. Alt



全国订货电话 4008-723-722

	ernative splicing results in multiple transcript variants. Re ad-through transcription also exists between this gene a nd the neighboring downstream C18orf32 (chromosome 18 open reading frame 32) gene.
Applications:	ELISA, WB, IHC
Name of antibody:	RPL17
Immunogen:	Fusion protein of human RPL17
Full name:	ribosomal protein L17
Synonyms:	L17; PD-1; RPL23
SwissProt:	P18621
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human tonsil
IHC Recommend dilution:	50-200
WB Predicted band size:	21 kDa
WB Positive control:	293T cell, Mouse brain tissue, Mouse testis tissue, HepG 2 cell, Mouse liver tissue, A375 cell lysates
WB Recommended dilution:	1000-5000





全国订货电话 4008-723-722

