

兔抗 NFKB2(Phospho-Ser866) 多克隆抗体

中文名称：兔抗 NFKB2(Phospho-Ser866) 多克隆抗体

英文名称：Anti-NFKB2(Phospho-Ser866) rabbit polyclonal antibody

别名：p52; p105; H2TF1; LYT10; CVID10; LYT-10; NF-kB2

抗原：NFKB2(Phospho-Ser866)

储存：冷冻（-20℃）避光

宿主：Rabbit

反应种属：Human Mouse Rat

相关类别：一抗

标记物：Unconjugate

克隆类型：Unconjugate

技术规格

Background:

NF-kappa-B is a pleiotropic transcription factor present in almost all cell types and is the endpoint of a series of signal transduction events that are initiated by a vast array of stimuli related to many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52. The dimers bind at kappa-B sites in the DNA of their target genes and the individual dimers have distinct preferences for different kappa-B sites that they can bind with distinguishable affinity and specificity. Different dimer combinations act as transcriptional activators or repressors, respectively.

Applications:	WB, IHC, IF
Name of antibody:	NFKB2(Phospho-Ser866)
Immunogen:	Peptide sequence around phosphorylation site of serine 866(E-D-S(p)-A-Y) derived from Human NFκB-p100/p52.
Full name:	nuclear factor of kappa light polypeptide gene enhancer in B-cells 2 (p49/p100)
Synonyms :	p52; p105; H2TF1; LYT10; CVID10; LYT-10; NF-κB2
SwissProt:	Q00653
IHC positive control:	Human breast carcinoma tissue
IHC Recommend dilution:	50-100
WB Predicted band size:	97 kDa
WB Positive control:	Human ovary cancer cells lysate
WB Recommended dilution:	500-1000
IF positive control:	Hela cells
IF Recommend dilution:	100-200



