

兔抗 NKD2 多克隆抗体

中文名称：兔抗 NKD2 多克隆抗体

英文名称：Anti-NKD2 rabbit polyclonal antibody

别名：naked cuticle homolog 2; Naked2

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：NKD2

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	This gene encodes a member of a family of proteins that function as negative regulators of Wnt receptor signaling through interaction with Dishevelled family members. The encoded protein participates in the delivery of transforming growth factor alpha-containing vesicles to the cell membrane. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.
Applications:	ELISA, WB, IHC
Name of antibody:	NKD2
Immunogen:	Fusion protein of human NKD2
Full name:	naked cuticle homolog 2
Synonyms:	Naked2
SwissProt:	Q969F2
ELISA Recommended dilution:	5000-10000

IHC positive control:	Human tonsil and Human prostate cancer
IHC Recommend dilution:	40-200
WB Predicted band size:	50 kDa
WB Positive control:	Mouse Pancreas tissue lysate
WB Recommended dilution:	500-2000



