

兔抗 PCYT2 多克隆抗体

| | |
|-------|--|
| 中文名称: | 兔抗 PCYT2 多克隆抗体 |
| 英文名称: | Anti-PCYT2 rabbit polyclonal antibody |
| 别名: | phosphate cytidyltransferase 2, ethanolamine; ET |
| 相关类别: | 一抗 |
| 储存: | 冷冻 (-20℃) |
| 宿主: | Rabbit |
| 抗原: | PCYT2 |
| 反应种属: | Human, Mouse, Rat |
| 标记物: | Unconjugate |
| 克隆类型: | rabbit polyclonal |

技术规格

| | |
|------------------------------------|---|
| Background: | This gene encodes an enzyme that catalyzes the formation of CDP-ethanolamine from CTP and phosphoethanolamine in the Kennedy pathway of phospholipid synthesis. Alternative splicing results in multiple transcript variants. |
| Applications: | ELISA, WB, IHC |
| Name of antibody: | PCYT2 |
| Immunogen: | Fusion protein of human PCYT2 |
| Full name: | phosphate cytidyltransferase 2, ethanolamine |
| Synonyms: | ET |
| SwissProt: | Q99447 |
| ELISA Recommended dilution: | 5000-10000 |
| IHC positive control: | Human cervical cancer and Human prostate cancer |

| | |
|---------------------------------|---|
| IHC Recommend dilution: | 40-200 |
| WB Predicted band size: | 44 kDa |
| WB Positive control: | Human fetal liver tissue, HepG2, A172, Hela and Jurkat cell lysates |
| WB Recommended dilution: | 1000-5000 |



