

兔抗 SLC10A7 多克隆抗体

中文名称: 兔抗 SLC10A7 多克隆抗体

英文名称: Anti-SLC10A7 rabbit polyclonal antibody

别 名: P7; C4orf13

相关类别: 一抗

储 存: 冷冻(-20℃) 避光

宿 主: Rabbit

抗 原: SLC10A7

反应种属: Human Mouse

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:

ti-pass membrane protein belonging to the sodiu m:bile acid symporter family. Existing as seven alt ernatively spliced isoforms, SLC10A7 is expressed at high levels in liver and lung, moderate levels in placenta, kidney, spleen, and thymus, and low I evels in heart, prostate, and testis. A few membe

rs of the sodium:bile acid symporter family, such as NTCP (also known as SLC10A1) and Asbt (also

known as SLC10A2), are involved in maintaining

The SLC10 family of sodium/bile salt cotransporte rs contains over 50 members in animal, plant an d bacterial species. SLC10A7 (solute carrier family 10 (sodium/bile acid cotransporter family), member 7), also known as P7, is a 358 amino acid mul



全国订货电话 4008-723-722

Applications:	enterohepatic circulation of bile acids by mediating the first step of active bile transport through membrane barriers of liver and intestine. Other family members, including SLC10A6, play an important role in the cellular delivery of specific prohormones in testis, placenta, adrenal gland and other peripheral tissues. Orphan carriers such as SLC10A7 are uncharacterized and their functions unknown.
Name of antibody:	SLC10A7
Immunogen:	Synthesized peptide derived from internal of hum an SLC10A7.
Full name:	solute carrier family 10, member 7
Synonyms:	P7; C4orf13
SwissProt:	Q0GE19
WB Predicted band size:	39 kDa
WB Positive control:	Mouse liver tissue lysate
WB Recommended dilution:	500-3000

