

兔抗 SLC22A6 多克隆抗体

中文名称：兔抗 SLC22A6 多克隆抗体

英文名称：Anti-SLC22A6 rabbit polyclonal antibody

别名：OAT1, PAHT, HOAT1, ROAT1

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：SLC22A6

反应种属：Human, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

The protein encoded by this gene is involved in the sodium-dependent transport and excretion of organic anions, some of which are potentially toxic. The encoded protein is an integral membrane protein and may be localized to the basolateral membrane. Four transcript variants encoding four different isoforms have been found for this gene. Functions as organic anion exchanger when the uptake of one molecule of organic anion is coupled with an efflux of one molecule of endogenous dicarboxylic acid (glutarate, ketoglutarate, etc). Mediates the sodium-independent uptake of 2,3-dimercapto-1-propanesulfonic acid (DMPS) By sim

| | |
|------------------------------------|--|
| | ilarity. |
| Applications: | ELISA, WB, IHC |
| Name of antibody: | SLC22A6 |
| Immunogen: | Synthetic peptide of human SLC22A6 |
| Full name: | Solute carrier family 22 (organic anion transporter), member 6 |
| Synonyms : | OAT1, PAHT, HOAT1, ROAT1 |
| SwissProt: | Q4U2R8 |
| ELISA Recommended dilution: | 2000-10000 |
| IHC positive control: | Human colon cancer and Human liver cancer |
| IHC Recommend dilution: | 50-200 |
| WB Predicted band size: | 56 kDa |
| WB Positive control: | Human fetal kidney tissue |
| WB Recommended dilution: | 500-2000 |



