

## 兔抗 PRKD3 多克隆抗体

中文名称: 兔抗 PRKD3 多克隆抗体

英文名称: Anti-PRKD3 rabbit polyclonal antibody

别名: EPK2; PKD3; PRKCN; PKC-NU; nPKC-NU

相关类别: 一抗

储存: 冷冻 (-20℃)

宿主: Rabbit

抗原: PRKD3

反应种属: Human, Mouse

标记物: Unconjugate

克隆类型: rabbit polyclonal

### 技术规格

|                          |   |
|--------------------------|---|
| <b>Background:</b>       | Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role. The protein encoded by this gene is one of the PKC family members. |
| <b>Applications:</b>     | ELISA, WB, IHC  |
| <b>Name of antibody:</b> | PRKD3   |
| <b>Immunogen:</b>        | Synthetic peptide of human PRKD3  |

|                                    |   |
|------------------------------------|---|
| <b>Full name:</b>                  | protein kinase D3   |
| <b>Synonyms :</b>                  | EPK2; PKD3; PRKCN; PKC-NU; nPKC-NU                            |
| <b>SwissProt:</b>                  | O94806  |
| <b>ELISA Recommended dilution:</b> | 1000-2000   |
| <b>IHC positive control:</b>       | Human cervical cancer and Human colon cancer                  |
| <b>IHC Recommend dilution:</b>     | 50-200  |
| <b>WB Predicted band size:</b>     | 100 kDa   |
| <b>WB Positive control:</b>        | Human testis tissue, K562, A549, Raji, NIH/3T3 and HeLa cells |
| <b>WB Recommended dilution:</b>    | 200-1000  |



