

兔抗 PROC 多克隆抗体

中文名称：兔抗 PROC 多克隆抗体

英文名称：Anti-PROC rabbit polyclonal antibody

别名：protein C, inactivator of coagulation factors Va and VIIIa; PC; APC; PROC1; THPH3; THPH4

相关类别：一抗

抗原：PROC

储存：冷冻（-20℃）

宿主：Rabbit

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	This gene encodes a vitamin K-dependent plasma glycoprotein. The encoded protein is cleaved to its activated form by the thrombin-thrombomodulin complex. This activated form contains a serine protease domain and functions in degradation of the activated forms of coagulation factors V and VIII. Mutations in this gene have been associated with thrombophilia due to protein C deficiency, neonatal purpura fulminans, and recurrent venous thrombosis.
Applications:	ELISA, WB, IHC
Name of antibody:	PROC
Immunogen:	Synthetic peptide of human PROC
Full name:	protein C, inactivator of coagulation factors Va and VIIIa
Synonyms:	PC; APC; PROC1; THPH3; THPH4

SwissProt:	P04070
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human prostate cancer and Human gastric cancer
IHC Recommend dilution:	50-100
WB Predicted band size:	52 kDa
WB Positive control:	SKOV3 cell lysate
WB Recommended dilution:	500-2000



