

兔抗 TACC3 多克隆抗体

中文名称：兔抗 TACC3 多克隆抗体

英文名称：Anti-TACC3 rabbit polyclonal antibody

别名：ERIC1; ERIC-1

相关类别：一抗

储存：冷冻（-20℃）避光

宿主：Rabbit

抗原：TACC3

反应种属：Human Mouse

标记物：Unconjugate

克隆类型：Unconjugate

技术规格

Background:	This gene encodes a member of the transforming acidic coiled-coil protein family. The encoded protein is a motor spindle protein that may play a role in stabilization of the mitotic spindle. This protein may also play a role in growth and differentiation of certain cancer cells.
Applications:	WB, IHC, IF
Name of antibody:	TACC3
Immunogen:	Synthesized peptide derived from C-terminal of human TACC3.
Full name:	transforming, acidic coiled-coil containing prote

	in 3
Synonyms :	ERIC1; ERIC-1
SwissProt:	Q9Y6A5
IHC positive control:	Human lung carcinoma tissue
IHC Recommend dilution:	50-100
WB Predicted band size:	90 kDa
WB Positive control:	293 cells and Huvec cells lysates
WB Recommended dilution:	500-3000
IF positive control:	A549cells
IF Recommend dilution:	100-500



