

兔抗 CCL17 多克隆抗体

中文名称: 兔抗 CCL17 多克隆抗体

英文名称: Anti-CCL17 rabbit polyclonal antibody

别名: TARC, ABCD-2, SCYA17, A-152E5.3

相关类别: 一抗

储存: 冷冻 (-20℃)

宿主: Rabbit

抗原: CCL17

反应种属: Human, Mouse

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:	This gene is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 16. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for T lymphocytes, but not monocytes or granulocytes. The product of this gene binds to chemokine receptors CCR4 and CCR8. This chemokine plays important roles in T cell development in thymus as well as in trafficking and activation of mature T cells.
Applications:	ELISA, WB, IHC
Name of antibody:	CCL17

Immunogen:	Synthetic peptide of human CCL17
Full name:	chemokine (C-C motif) ligand 17
Synonyms :	TARC, ABCD-2, SCYA17, A-152E5.3
SwissProt:	Q92583
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human colon cancer and Human gastric cancer
IHC Recommend dilution:	100-300
WB Predicted band size:	11 kDa
WB Positive control:	Mouse lung and thymus tissue, human fetal muscle tissue
WB Recommended dilution:	500-2000



